

A short proof of a formula of Brasselet, Lê and Seade for the Euler obstruction

Jörg Schürmann

Westf. Wilhelms-Universität, SFB 478

”Geometrische Strukturen in der Mathematik”

Hittorfstr. 27, 48149 Münster, Germany

E-mail: jschuerm@math.uni-muenster.de

Using the index theorem of Dubson, Lê, Ginsburg and Sabbah for the vanishing cycle functor (which is a special case of [Sch2]), we give a short proof of a generalization of the following theorem [BLS, thm.3.1]:

Theorem 0.1. *Let $(X, 0)$ be a germ of an equidimensional complex space (of positive dimension) in $\mathbb{C}^N$. Let V_i , $i = 1, \dots, k$, be the (connected) strata of a Whitney stratification of a small representative X of $(X, 0)$. Then there is an open dense Zariski subset Ω in the space of complex linear forms $\mathbb{L}$ on $\mathbb{C}^N$, such that for every $l \in \Omega$, there is ϵ_0 , such that for any ϵ , $\epsilon_0 > \epsilon > 0$ and t_0 sufficiently small, we have the following formula for the Euler obstruction of $(X, 0)$:*

$$Eu(X, 0) = \sum_1^k \chi(V_i \cap \mathbb{B}_\epsilon \cap l^{-1}(t_0)) \cdot Eu_{V_i}(X),$$

where χ denotes the Euler-Poincaré characteristic and $Eu_{V_i}(X)$ is the value of the Euler obstruction of X at any point of V_i , $i = 1, \dots, k$.

Here $\mathbb{B}_\epsilon$ is the corresponding closed ball in $\mathbb{C}^N$, and χ is the usual Euler characteristic

$$\chi(Y) := \chi(H^*(Y, \mathbb{Q}))$$

of a topological space Y with finite dimensional cohomology $H^*(Y, \mathbb{Q})$. Similarly, let us denote by $\chi_c(Y)$ the Euler characteristic of $H_c^*(Y, \mathbb{Q})$ for locally compact Y (if defined).

Remark 0.1. *This result goes already back to an old preprint [Du] of Dubson (compare with [Du, thm. V.1.1, p.69] (Réduction à la section hyperplane) and [Du, thm. IV.4.1.2, p.67] (Lemme de Hopf singulier)), which was not mentioned in [BLS].*

Consider a Whitney stratification of a germ of a complex space $(X, 0)$ of positive dimension in $\mathbb{C}^N$ as above, but this time we don't assume $(X, 0)$ is equidimensional. Let us refine the stratification by adding the point stratum $V_0 := \{0\}$, and then assume $0 \notin V_i$ for $i > 0$. Take a linear form $l \in \mathbb{L}$, such that dl_0 is a nondegenerate covector in the sense of stratified Morse theory (i.e. it does not vanish on any generalized tangent space $\lim_{x_n \rightarrow 0} T_{x_n} V_i$ at 0, with x_n a sequence in V_i , $i = 1, \dots, k$). This condition defines a Zariski open dense subset Ω of $\mathbb{L}$. Then

$$V(i, \epsilon, t_0) := V_i \cap \mathbb{B}_\epsilon \cap l^{-1}(t_0)$$

is for $i > 0$ a smooth manifold with boundary for $0 < |t_0| \ll \epsilon_0 \ll 1$ (this is also true for any $l \in \mathbb{L}$, since one can refine our stratification to an a_f -stratification). By Poincaré-duality we get

$$\chi(V(i, \epsilon, t_0)) = (-1)^{2(\dim_{\mathbb{C}}(V_i)-1)} \cdot \chi_c(V^\circ(i, \epsilon, t_0)) = \chi_c(V^\circ(i, \epsilon, t_0)), \quad (1)$$

with $V^\circ(i, \epsilon, t_0) := V_i \cap \mathbb{B}_\epsilon^\circ \cap l^{-1}(t_0)$, and $\mathbb{B}_\epsilon^\circ$ the corresponding open ball.

Consider now a function $\alpha : X \rightarrow \mathbb{Z}$, which is constructible with respect to the refined Whitney stratification (i.e. is constant on the strata). Let $i_0 : \{0\} \rightarrow X \cap \{l = 0\}$ and $i : X \cap \{l = 0\} \rightarrow X$ be the inclusions. Then we have for $0 < |t_0| \ll \epsilon_0 \ll 1$:

$$i_0^!(\psi_l(\alpha))(0) = \sum_1^k \chi_c(V^\circ(i, \epsilon, t_0)) \cdot \alpha_i, \quad (2)$$

with α_i the value of α on V_i , and ψ_l the nearby cycle functor with respect to l on the level of constructible functions (and similarly for $i_0^!$). This can be defined by two different (but equivalent) approaches:

- One can use the sophisticated language of analytically constructible complexes of sheaves:

Choose a bounded complex $\mathcal{F}$ of sheaves of $\mathbb{C}$ -vector spaces on X , which is constructible with respect to our refined Whitney stratification, such that all stalk complexes $\mathcal{F}_x$ have finite dimensional cohomology with $\chi(H^*(\mathcal{F}_x)) = \alpha(x)$ for all $x \in X$. Then

$$i_0^!(\psi_l(\alpha))(0) := \chi(H^*(i_0^! \psi_l(\mathcal{F}))). \quad (3)$$

This is well defined by [Sch, Part II.3, Part VI], and equation (2) follows from [Sch, Part V, lem.3.4].

- By linearity one can assume $\alpha = 1_Z$, with Z a closed union of strata and then one uses a direct geometric definition in terms of a corresponding local Milnor fibration:

$$i_0^!(\psi_l(1_Z))(0) = \chi_c(M^\circ(l|Z)) = \sum_{V_i \subset Z} \chi_c(V^\circ(i, \epsilon, t_0)), \quad (4)$$

with $M^\circ(l|Z) := Z \cap \mathbb{B}_\epsilon^\circ \cap l^{-1}(t_0)$ (for $0 < |t_0| \ll \epsilon_0 \ll 1$) the open local Milnor fiber of $l|Z$ at 0.

But by [Sch, Part VI, ex.2.3] we have

$$i_0^!(\psi_l(\alpha))(0) = i_0^*(\psi_l(\alpha))(0). \quad (5)$$

In our situation this formula can also be explained more geometrically in the following way: Choose a Whitney stratification adapted to α and $\{l = 0\}$ (e.g. $\{l = 0\}$ is a union of strata), which is also an a_f -stratification. Then the corresponding local Milnor fibrations are "locally constant" along the strata of $\{l = 0\}$ so that $\psi_l(\alpha)$ is constructible with respect to the induced stratification of $\{l = 0\}$. So by linearity it is again enough to show

$$i_0^!(1_Z)(0) = i_0^*(1_Z)(0) \quad (6)$$

for Z a closed union of strata of $\{l = 0\}$. But $i_0^*(1_Z)(0) - i_0^!(1_Z)(0)$ is by the locally conic structure of Z just the Euler characteristic of the link $Z \cap \{|z| = \epsilon\}$ of Z in 0 (for $0 < \epsilon \ll 1$), which vanishes by a classical result of Sullivan (compare [BV, Su] and also [KS, Ex.IX.12, p.409]).

From the definition of the vanishing cycle functor, we get the equality

$$\alpha(0) = i_0^* i^*(\alpha)(0) = i_0^*(\psi_l(\alpha))(0), \quad (7)$$

if $\chi(H^*(i_0^* \phi_l(\mathcal{F}))) = 0$. Or more directly in terms of constructible functions:

$$i_0^*(\phi_l(\alpha))(0) := i_0^*(\psi_l(\alpha))(0) - i_0^* i^*(\alpha)(0).$$

But this follows from the index theorem of Dubson, Lê, Ginsburg and Sabbah [Du2, Lê, Lê2, Gi, Sa] for the vanishing cycle functor, if the graph of dl doesn't intersect the support $|CC(\mathcal{F})| = |CC(\alpha)|$ of the characteristic cycle of $\mathcal{F}$ or α in $T^*\mathbb{C}^N$ over an open neighborhood U of 0. More precisely, this index formula implies ([Sa, thm.4.5, rem.4.6]):

$$-i_0^*(\phi_l(\alpha))(0) = -\chi(H^*(i_0^* \phi_l(\mathcal{F}))) = \sharp(dl(U) \cap CC(\mathcal{F})), \quad (8)$$

if $dl(U) \cap |CC(\mathcal{F})| \subset \{(0, dl_0)\}$.

Remark 0.2. For another purely real proof of this intersection formula compare with [Sch2] (which is also closely related to [KS, thm.9.5.6]).

Let $Eu(cl(V_i), \cdot)$ be the Euler obstruction of the closure of the stratum V_i ($i > 0$). This function is constructible with respect to our refined Whitney stratification (compare [Sch, part VI] or [CEP, exp.6, cor.10.2, p.125]), with

$$|CC(Eu(cl(V_i), \cdot))| = cl(T_{V_i}^* \mathbb{C}^N).$$

But the condition $(0, dl_0) \notin cl(T_{V_i}^* \mathbb{C}^N)$ is equivalent to the fact, that dl_0 does not vanish on any generalized tangent space $\lim_{x_n \rightarrow 0} T_{x_n} V_i$ at 0, with x_n a sequence in V_i ($i \in \{1, \dots, k\}$).

Altogether we get the following generalization of theorem 0.1 of Brasselet, Lê and Seade:

Theorem 0.2. Let $(X, 0)$ be a germ of a complex space (of positive dimension) in $\mathbb{C}^N$. Let V_i , $i = 1, \dots, k$, be the (connected) strata of a Whitney stratification of a small representative X of $(X, 0)$. Refine the stratification by adding the point stratum $V_0 := \{0\}$, and then assume $0 \notin V_i$ for $i > 0$. Fix a (constructible) function α , which is a linear combination of $Eu(cl(V_i), \cdot)$ (for some $i \in I \subset \{1, \dots, k\}$). Consider a complex linearform $l \in \mathbb{L}$, with dl_0 not vanishing on any generalized tangent space $\lim_{x_n \rightarrow 0} T_{x_n} V_i$ at 0, with x_n a sequence in V_i ($i \in I$). Then there is an $\epsilon_0 > 0$ with

$$\alpha(0) = \sum_1^k \chi_c(V^\circ(i, \epsilon, t_0)) \cdot \alpha_i = \sum_1^k \chi(V_i \cap \mathbb{B}_\epsilon \cap l^{-1}(t_0)) \cdot \alpha_i,$$

for $0 < |t_0| < \epsilon < \epsilon_0$.

Remark 0.3. The same result is true with the same proof, if the stratification satisfies suitable weaker regularity conditions than Whitney b -regularity, e.g. the stratification is (locally) C -regular in the sense of Bekka [Be] (compare with [Sch, Part VI], even for a more general class of stratifications). Moreover, this formula is more generally true (with the same proof) for any holomorphic function germ $l : (\mathbb{C}^N, 0) \rightarrow (\mathbb{C}, 0)$, with dl_0 as before.

References

- [Be] Bekka.K.: C -régularité et trivialité topologique. in: Warwick Singularity Symposium (1989), Lect. Notes in Math. 1462, 42-62 (1991)

- [BLS] Brasselet, J.P., Lê D.T., Seade, J.: Euler obstruction and indices of vector fields. *Topology* 39, 1193-1208 (2000)
- [BV] Burghela, D., Verona, V.: Local homological properties of analytic sets. *Manus. math.* 7, 55-66 (1972)
- [CEP] Caractéristique d'Euler-Poincaré. Séminaire E.N.S. 1978-1979. *Astérisque* 82-83 (1981)
- [Du] Dubson, A.: Calcul des invariants numériques des singularités at applications. preprint, Sonderforschungsbereich 40, Univ. of Bonn (1981)
- [Du2] Dubson, A.: Formule pour l'indice des complexes constructibles et D -modules holonomes. *C. R. Acad. Sci. Paris* 298, 113-164 (1984)
- [Gi] Ginsburg, V.: Characteristic cycles and vanishing cycles. *Inv. Math.* 84, 327-402 (1986)
- [KS] Kashiwara, M., Schapira, P.: *Sheaves on Manifolds*. Springer, Berlin Heidelberg (1990)
- [Lê] Lê D.T.: Le concept de singularité isolée de fonction analytique. *Advanced Studies in Pure Math.* 8, 215-227 (1986)
- [Lê2] Lê D.T.: Morsification of D -Modules. *Bol. Soc. Mat. Mexicana* 3, 229-248 (1998)
- [Sa] Sabbah, C.: Quelques remarques sur la géométrie des espaces conormaux. *Astérisque*. 130, 161-192 (1985)
- [Sch] Schürmann, J.: Topology of singular spaces and constructible sheaves. Habilitation-paper, 325 pp, Hamburg (2000), available at <http://www.math.uni-hamburg.de/home/schuermann>
- [Sch2] Schürmann, J.: A general intersection formula for Lagrangian cycles. preprint, math.AG/0201300
- [Su] Sullivan, D.: Combinatorial invariants of analytic spaces. *Proc. of Liverpool singularities symposium I, Lect. Notes in Math.* 192, 165-168 (1971)